CLINICAL REFERENCE DOSSIER

The Patient's Guide to CO_2 Laser Resurfacing

A medical analysis of ablative and fractional laser neocollagenesis, day-by-day healing stages, and candidacy for deep facial tissue restoration.

Dr Bradley Tomkins

MBChB, MRCGP • GMC Registered Medical Practitioner

Former Royal Navy Medical Officer • Laser & Aesthetic Physician
Doctor Brad Laser & Cosmetic Clinic, Fishponds, Bristol

1. THE MECHANISM OF ACTION: DEEP DERMAL NEOCOLLAGENESIS

The human epidermis is engineered as an impermeable environmental shield. While non-medical treatments (such as chemical peels, hydradermabrasion, and shallow microneedling) can refine superficial dead keratinocytes, they cannot reach the reticular dermis where structural skin slackness and etched wrinkles originate.

The REVIDA carbon dioxide (CO_2) laser emits coherent light at a wavelength of **10,600 nanometers**, targeting intracellular water within damaged cutaneous tissue. The clinical impact occurs across two simultaneous physiological phases:

> **Phase 1: Controlled Micro-Ablation**
> Targeted columns of photodamaged epidermis and solar elastosis are vaporized instantly to precise microscopic depths, eliminating surface irregularity and cellular atypia.

> **Phase 2: Thermal Neocollagenesis (65°C – 70°C)**
> Surrounding zones receive controlled lateral thermal diffusion. This heat denatures degenerate collagen fibers, causing immediate tissue tightening, while signaling fibroblast activation to produce fresh **Type-I and Type-III collagen** over 3 to 6 months.

2. CLINICAL PAIN MANAGEMENT: LOCAL & REGIONAL NERVE BLOCKS

Fear of operative pain is the most frequent deterrent for patients considering ablative laser therapy. In commercial salons, non-medical practitioners are legally restricted to over-the-counter topical creams (such as EMLA or 4% lidocaine), which provide minimal numbing for deep thermal energy.

As a medical doctor, my protocol eliminates sharp pain through clinical regional anesthesia:

Anesthesia Modality	Anatomical Target	Clinical Benefit to Patient
Regional Dental Blocks	Infraorbital & Mental sensory nerves	Completely desensitizes upper/lower lips and perioral zones. Completely eliminates sharp laser sensation.
Supraorbital Blocks	Forehead & Glabella nerve branches	Provides profound numbing for deep brow wrinkles and forehead resurfacing without sedation.
Subdermal Infiltration	Localized delicate zones & crows feet	Pinpoint targeted lidocaine numbing for focused periorbital passes.
Concomitant Chilling	Epidermal surface airflow (-20°C air)	Cools the skin during energy delivery, reducing thermal discomfort and post-procedure heat.

With regional blocks active, the treatment is calm and well tolerated. Patients experience mild vibrational pressure and radiant warmth rather than sharp pain.

3. THE DAY-BY-DAY BIOLOGICAL HEALING TIMELINE

Downtime is not an unwanted side effect; it is the biological process of tissue replacement. Because REVIDA physically removes aged skin, the body requires a structured recovery window to complete re-epithelialization. Plan for **7 to 10 days of social recovery** for full-field or deep fractional protocols.

Day 0 (Procedure Day): The Immediate Response
The treated skin appears deeply flushed, similar to moderate sunburn, and feels hot. A medical occlusive barrier (such as sterile white soft paraffin or Aquaphor) is applied before leaving the clinic. Discomfort is managed with standard paracetamol or ibuprofen.

Days 1 to 2: Peak Edema & Serous Exudate
Swelling peaks, especially around the eyes upon waking. The micro-ablative columns weep clear serous fluid. Patients perform gentle, dilute sterile vinegar soaks (which maintain skin acidity and prevent bacterial colonization) followed by immediate reapplication of occlusive barrier balm 4 to 5 times daily.

Days 3 to 4: The Shedding & "Sandpaper" Phase
Serous weeping ceases. The ablated skin turns a bronzed, light brown shade and feels rough and dry, like fine sandpaper. Micro-crusting begins sloughing away from the center of the face outwards during cleansing. *Crucial rule:* Skin must never be picked or manually peeled; it must shed passively.

Days 5 to 7: Re-Epithelialization Complete
The bronzed outer layer sloughs off completely. Beneath it sits newly synthesized, fresh, pink, velvety skin. Facial edema has resolved. Most patients feel comfortable resuming non-strenuous desk work and light social routines at this stage.

Days 8 to 14: Erythema Fade & Daily Photoprotection
The intense pinkness gradually softens into a mild flush. Mineral foundation or tinted SPF 50+ may be applied. Strict broad-spectrum UV protection is essential to shield vulnerable young melanocytes from hyperpigmentation.

Months 1 to 6: Long-Term Neocollagenesis
While surface skin is completely healed by day 10, deep thermal collagen remodeling continues for up to 6 months, yielding progressive tightening of perioral lines, pore reduction, and structural skin density.

4. CLINICAL CANDIDACY MATRIX: SUITABILITY & SAFETY

Patient safety requires honest assessment. CO_2 laser energy is powerful; treating inappropriate skin types or unmanaged medical conditions risks complications such as post-inflammatory hyperpigmentation (PIH) or scarring.

Clinical Indication	Candidacy Status	Medical Rationale
Fitzpatrick Skin Types I – III (Fair to light-olive European skin)	**Optimal Candidate**	Low baseline epidermal melanin. Minimal risk of post-inflammatory hyperpigmentation under proper sun avoidance.
Fitzpatrick Skin Types IV – VI (Darker olive, Asian, or Black skin)	**Contraindicated for Ablative CO_2**	High risk of permanent hyper- or hypopigmentation. Alternative non-ablative or pigment laser wavelengths are recommended.
Perioral Lines ("Smoker's Lines")	**Prime Indication**	Dermal thinning around the mouth responds well to confluent CO_2 ablation and structural collagen contraction.
Roaccutane / Isotretinoin Use	**Contraindicated**	Must have stopped oral isotretinoin for at least 6 to 12 months prior to ensure normal follicular wound healing.

5. THE DIAGNOSTIC MEDICAL ASSESSMENT (£60)

Every laser procedure, parameter setting, and follow-up at our Bristol practice is conducted personally by Dr Bradley Tomkins. We do not use sales coordinators or hurried 10-minute turnaround slots.

Your 30-minute clinical consultation includes a comprehensive facial assessment, skin thickness evaluation, downtime mapping against your diary, and a clear, fixed-fee quotation with zero pressure to book.

Ready to Explore REVIDA CO_2 Resurfacing?

Consultations are held at our private Fishponds clinic in Bristol. Check real-time diary availability online or send a direct clinical enquiry.

Reserve Consultation with Dr Brad (£60) →

Clinic WhatsApp: 07588 971425 • 583 Fishponds Road, Bristol, BS16 3AA